

What to know about **TECENTRIQ + Avastin®** (bevacizumab)

About **TECENTRIQ**

TECENTRIQ is a type of treatment known as immunotherapy. In HCC, cancer cells can hide from your immune system, which means the immune system can't attack the cancer cells like it normally would. Immunotherapy, like TECENTRIQ, is designed to help the immune system find and attack the cancer cells that are hiding. TECENTRIQ may also affect healthy cells. Many patients confuse immunotherapy with traditional chemotherapy. Immunotherapy is **not traditional chemotherapy**.

TECENTRIQ + Avastin: Proven results

Avastin is another treatment that is used with TECENTRIQ to treat hepatocellular carcinoma (HCC). Avastin is not traditional chemotherapy. TECENTRIQ plus Avastin was studied in a clinical trial against sorafenib, an oral medicine (pill) for HCC. The study included 501 adult patients with HCC that has spread or tumors that could not be removed by surgery.



More patients
LIVED LONGER

The study found that TECENTRIQ + Avastin lowered the risk of death by **42%**, compared to sorafenib.



LOWER CHANCE
of cancer spreading, growing,
or getting worse

Patients who received TECENTRIQ + Avastin were **41%** less likely to have their cancer get worse, compared to sorafenib.

How is **TECENTRIQ + Avastin** given?

Your healthcare team will give TECENTRIQ + Avastin with an intravenous or "IV" line. This means a bag of each medicine is given in your vein, one after the other. TECENTRIQ + Avastin may be given **every 3 weeks**. Your doctor may choose to give TECENTRIQ on a different schedule, so that you receive TECENTRIQ infusions every 2, 3, or 4 weeks. Your healthcare team will also decide **how long you may be on treatment**. If your disease gets worse or treatments can't be tolerated, your healthcare team may stop infusions.

First infusion

The first **TECENTRIQ** infusion is given over **60 minutes**.



TECENTRIQ



60 minutes

Following infusions

Depending on your response to the first infusion, later infusions of **TECENTRIQ** may take **30 minutes**.



TECENTRIQ



30 minutes

**More time will be needed for your other medications, including Avastin.
Ask your healthcare team how long the entire process will take.**

What is **TECENTRIQ**?

TECENTRIQ is a prescription medicine used to treat:

Adults with a type of liver cancer called hepatocellular carcinoma (HCC). TECENTRIQ may be used with the medicine bevacizumab when your liver cancer:

- has spread or cannot be removed by surgery, **and**
- you have not received other medicines by mouth or injection through your vein (IV) to treat your cancer.

It is not known if TECENTRIQ is safe and effective when used:

- in children for the treatment of HCC.

Important Safety Information

What is the most important information about TECENTRIQ?

TECENTRIQ can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during your treatment or even after your treatment has ended.

Please see additional Important Safety Information on the following page and accompanying full Prescribing Information.

 **TECENTRIQ®**
atezolizumab 840 mg / 1200 mg
INJECTION FOR IV USE

Important Safety Information (Continued)

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

Lung problems

- cough
- shortness of breath
- chest pain

Intestinal problems

- diarrhea (loose stools) or more frequent bowel movements than usual
- stools that are black, tarry, sticky, or have blood or mucus
- severe stomach-area (abdomen) pain or tenderness

Liver problems

- yellowing of your skin or the whites of your eyes
- severe nausea or vomiting
- pain on the right side of your stomach area (abdomen)
- dark urine (tea colored)
- bleeding or bruising more easily than normal

Hormone gland problems

- headaches that will not go away or unusual headaches
- eye sensitivity to light
- eye problems
- rapid heartbeat
- increased sweating
- extreme tiredness
- weight gain or weight loss
- feeling more hungry or thirsty than usual
- urinating more often than usual
- hair loss
- feeling cold
- constipation
- your voice gets deeper
- dizziness or fainting
- changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness

Kidney problems

- decrease in your amount of urine
- blood in your urine
- swelling of your ankles
- loss of appetite

Skin problems

- rash
- itching
- skin blistering or peeling
- painful sores or ulcers in mouth or nose, throat, or genital area
- fever or flu-like symptoms
- swollen lymph nodes

Problems can also happen in other organs.

These are not all of the signs and symptoms of immune system problems that can happen with TECENTRIQ. Call or see your healthcare provider right away for any new or worse signs or symptoms, including:

- Chest pain, irregular heartbeat, shortness of breath, or swelling of ankles
- Confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight
- Persistent or severe muscle pain or weakness, muscle cramps
- Low red blood cells, bruising

Infusion reactions that can sometimes be severe or life-threatening. Signs and symptoms of infusion reactions may include:

- chills or shaking
- dizziness

- itching or rash
- flushing
- shortness of breath or wheezing
- feeling like passing out
- fever
- back or neck pain

Complications, including graft-versus-host disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic). These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with TECENTRIQ. Your healthcare provider will monitor you for these complications.

Getting medical treatment right away may help keep these problems from becoming more serious. Your healthcare provider will check you for these problems during your treatment with TECENTRIQ. Your healthcare provider may treat you with corticosteroid or hormone replacement medicines. Your healthcare provider may also need to delay or completely stop treatment with TECENTRIQ if you have severe side effects.

Before you receive TECENTRIQ, tell your healthcare provider about all of your medical conditions, including if you:

- have immune system problems such as Crohn's disease, ulcerative colitis, or lupus
- have received an organ transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have received radiation treatment to your chest area
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. TECENTRIQ can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with TECENTRIQ. **Females who are able to become pregnant:**
 - Your healthcare provider should do a pregnancy test before you start treatment with TECENTRIQ.
 - You should use an effective method of birth control during your treatment and for at least 5 months after the last dose of TECENTRIQ.
- are breastfeeding or plan to breastfeed. It is not known if TECENTRIQ passes into your breast milk. Do not breastfeed during treatment and for at least 5 months after the last dose of TECENTRIQ.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

The most common side effects of TECENTRIQ when used in hepatocellular carcinoma (HCC) with bevacizumab include:

- high blood pressure
- feeling tired or weak
- too much protein in the urine

TECENTRIQ may cause fertility problems in females, which may affect the ability to have children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of TECENTRIQ. Ask your healthcare provider or pharmacist for more information about the benefits and side effects of TECENTRIQ.

You may report side effects to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. You may also report side effects to Genentech at 1-888-835-2555.

Please see accompanying full Prescribing Information and Medication Guide for additional Important Safety Information.

Genentech

A Member of the Roche Group

© 2023 Genentech USA, Inc. All rights reserved. M-US-00016115(v2.0)

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1200 mg
INJECTION FOR IV USE