


Preparing to talk to your doctor about TECENTRIQ

Your doctor has prescribed TECENTRIQ, an immunotherapy for adults with muscle-invasive bladder cancer (MIBC) who have something called circulating tumor DNA molecular residual disease (ctDNA MRD). A positive ctDNA MRD test means that small pieces of the tumor's DNA were detected in your blood, and the cancer may still be present in your body. You will receive TECENTRIQ after your bladder has been removed by surgery (cystectomy).

Your doctor is here to guide you on next steps. Be sure to share any questions and concerns with them so they can better support your care. This guide may be helpful in preparing for your next visit.

4 ways to prepare for your next doctor's appointment



Keep a treatment journal

so you and your caregiver can write down questions, concerns, and helpful information from your doctor



Ask a friend or family member to come

to help take notes and provide emotional support



Have your health information handy

including your health history, current medications, and insurance card



Advocate for yourself

by taking the time to ask questions, and by being open and honest with your doctor

WHAT IS TECENTRIQ?

TECENTRIQ is a prescription medicine used to treat:

Adults with a type of bladder cancer called muscle invasive bladder cancer (MIBC) that has spread into the muscle layer of the bladder but not to other parts of the body. TECENTRIQ may be used alone as a treatment for your bladder cancer:

- to help prevent your bladder cancer from coming back after your bladder has been removed by surgery, **and**
- small pieces of DNA from the tumor (called circulating tumor DNA [ctDNA]) were found in your blood, showing that cancer cells remain in the body (molecular residual disease). Your healthcare provider will perform a test to make sure that TECENTRIQ is right for you.

It is not known if TECENTRIQ is safe and effective when used:

- in children for the treatment of MIBC.

SELECT IMPORTANT SAFETY INFORMATION

TECENTRIQ can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during your treatment or even after your treatment has ended.

Please see full [Prescribing Information](#) and [Medication Guide](#) for additional Important Safety Information.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1200 mg
INJECTION FOR IV USE

Possible questions to ask your doctor

As you begin your treatment journey with TECENTRIQ, you may have some questions. Asking your doctor some of the following questions can help you get the answers you need to feel ready for every step.



Starting TECENTRIQ

- ☐ What are the potential benefits of receiving TECENTRIQ after surgery?
- ☐ What can TECENTRIQ do for me?
- ☐ What are the possible side effects of TECENTRIQ?
- ☐ How long will I be on treatment?



Receiving TECENTRIQ

- ☐ What options do I have to receive treatment with TECENTRIQ?
- ☐ Depending on how I receive TECENTRIQ, where will my treatment take place?
- ☐ How should I prepare for my first treatment?
- ☐ Can you walk me through a typical treatment day?



Managing your day-to-day during treatment

- ☐ How will this treatment affect my daily life?
- ☐ What should I do if I experience side effects?
- ☐ How will treatment impact my lifestyle?



Learn more about
financial support at
[TECENTRIQ.com](https://www.tecentriq.com)



Please see full [Prescribing Information](#) and [Medication Guide](#) for additional Important Safety Information.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1200 mg
INJECTION FOR IV USE

IMPORTANT SAFETY INFORMATION

What is the most important information about TECENTRIQ?

TECENTRIQ can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during your treatment or even after your treatment has ended.

Call or see your healthcare provider right away if you develop any new or worsening signs or symptoms, including:

Lung problems

- cough
- shortness of breath
- chest pain

Intestinal problems

- diarrhea (loose stools) or more frequent bowel movements than usual
- stools that are black, tarry, sticky, or have blood or mucus
- severe stomach-area (abdomen) pain or tenderness

Liver problems

- yellowing of your skin or the whites of your eyes
- severe nausea or vomiting
- pain on the right side of your stomach area (abdomen)
- dark urine (tea colored)
- bleeding or bruising more easily than normal

Hormone gland problems

- headaches that will not go away or unusual headaches
- eye sensitivity to light
- eye problems
- rapid heartbeat
- increased sweating
- extreme tiredness
- weight gain or weight loss
- feeling more hungry or thirsty than usual
- urinating more often than usual
- hair loss
- feeling cold
- constipation
- your voice gets deeper
- dizziness or fainting
- changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness

Kidney problems

- decrease in your amount of urine
- blood in your urine
- swelling of your ankles
- loss of appetite

Skin problems

- rash
- itching
- skin blistering or peeling
- painful sores or ulcers in your mouth or your nose, throat, or genital area
- fever or flu-like symptoms
- swollen lymph nodes

Problems can also happen in other organs.

These are not all of the signs and symptoms of immune system problems that can happen with TECENTRIQ. Call or see your healthcare provider right away for any new or worsening signs or symptoms, including:

- Chest pain, irregular heartbeat, shortness of breath, or swelling of ankles
- Confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight
- Persistent or severe muscle pain or weakness, muscle cramps
- Low red blood cells, bruising

Please see full [Prescribing Information](#) and [Medication Guide](#) for additional Important Safety Information.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1200 mg
INJECTION FOR IV USE

IMPORTANT SAFETY INFORMATION (cont'd)

Infusion reactions that can sometimes be severe or life-threatening. Signs and symptoms of infusion reactions may include:

- chills or shaking
- itching or rash
- flushing
- shortness of breath or wheezing
- dizziness
- feeling like passing out
- fever
- back or neck pain

Rejection of a transplanted organ or tissue. Your healthcare provider should tell you what signs and symptoms you should report and monitor you depending on the type of organ or tissue transplant that you have had.

Complications, including graft-versus-host disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic). These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with TECENTRIQ. Your healthcare provider will monitor you for these complications.

Getting medical treatment right away may help keep these problems from becoming more serious. Your healthcare provider will check you for these problems during your treatment with TECENTRIQ. Your healthcare provider may treat you with corticosteroid or hormone replacement medicines. Your healthcare provider may also need to delay or completely stop treatment with TECENTRIQ if you have severe side effects.

Before you receive TECENTRIQ, tell your healthcare provider about all of your medical conditions, including if you:

- have immune system problems such as Crohn's disease, ulcerative colitis, or lupus
- have received an organ or tissue transplant, including corneal transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have received radiation treatment to your chest area
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. TECENTRIQ can harm your unborn baby. Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with TECENTRIQ. **Females who are able to become pregnant:**
 - Your healthcare provider should do a pregnancy test before you start treatment with TECENTRIQ.
 - You should use an effective method of birth control during your treatment and for at least 5 months after the last dose of TECENTRIQ.
- are breastfeeding or plan to breastfeed. It is not known if TECENTRIQ passes into your breast milk. Do not breastfeed during treatment and for at least 5 months after the last dose of TECENTRIQ.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

The most common side effects of TECENTRIQ when used alone include:

- feeling tired or weak
- decreased appetite
- nausea
- cough
- shortness of breath

TECENTRIQ may cause fertility problems in females, which may affect the ability to have children. Talk to your healthcare provider if you have concerns about fertility.

These are not all the possible side effects of TECENTRIQ. Ask your healthcare provider or pharmacist for more information about the benefits and side effects of TECENTRIQ.

You may report side effects to the FDA at [1-800-FDA-1088](tel:1-800-FDA-1088) or www.fda.gov/medwatch.
You may also report side effects to Genentech at [1-888-835-2555](tel:1-888-835-2555).

Please see full [Prescribing Information](#) and [Medication Guide](#) for additional Important Safety Information.

 **TECENTRIQ**[®]
atezolizumab 840 mg / 1200 mg
INJECTION FOR IV USE